

---

# First record of Great Knot *Calidris tenuirostris* for Namibia

Josh Engel

---

**Première mention du Bécasseau de l'Anadyr *Calidris tenuirostris* pour la Namibie.** Un Bécasseau de l'Anadyr *Calidris tenuirostris* a été observé et photographié sur la lagune de Walvis Bay, en Namibie, le 24 novembre 2008. Ceci constitue la première donnée pour le pays. L'aire d'hivernage de cette espèce, qui niche au nord-est de la Sibérie, s'étend de l'Australasie jusqu'au Pakistan. En Afrique, des occasionnels ont été notés au Maroc (un individu en août 1980), au Mozambique (un en décembre 2004 ; trois en septembre 2008) et sur la côte ouest de l'Afrique du Sud (un, supposé être le même individu, en quatre saisons consécutives en 2000–03).

On the afternoon of 24 November 2008, I was birding with my tour group at the Walvis Bay Lagoon on the central Namibian coast. The lagoon's shoreline was full of waders on the outgoing tide, with large numbers of Little Stints *Calidris minuta*, Curlew Sandpipers *C. ferruginea* and Sanderlings *C. alba* on the exposed mudflats. Larger waders were slightly further out, at the tideline, including many Bar-tailed Godwits *Limosa lapponica* and Whimbrels *Numenius phaeopus*.

As I was scanning the flocks, I picked out a different-looking wader amongst the Bar-tailed Godwits. It was far smaller than the godwits, but larger than the *Calidris* sandpipers, with a fairly long, hefty, slightly decurved bill (Figs. 1–2). I immediately recognised it as a Great Knot *Calidris tenuirostris*, a species I am familiar with from having observed thousands and ringed hundreds in

Australia. Whilst superficially similar to Red Knot *C. canutus*, Great Knot is a larger bird, with a longer bill, a less contrasting facial pattern and a white rump, seen while the bird bathed.

We watched and photographed the bird for the next 30 minutes at a distance of c.40 m. At one point it took off with a group of other waders, but we quickly relocated it. While it was originally feeding with the larger waders, after it flew it joined the smaller species. It also bathed, preened and rested during our brief period of observation. Unfortunately I was unable to age the bird in the field and the photographs are inconclusive.

Great Knot is largely restricted to the East Asian-Australasian flyway, breeding in northern Siberia and spending the non-breeding season as far west as coastal Pakistan (Hayman *et al.* 1986). It is now known to winter as far west as the east and south-east coasts of the Arabian Peninsula (Aspinall 1994, Green *et al.* 1994).

Whilst Great Knot was first recorded on the African continent in Morocco in August 1980 (Lister 1981), it has only been recorded in sub-Saharan Africa in the last decade (Cohen & Winter 2003). The Walvis Bay bird represents the first record for Namibia. Hockey *et al.* (2005) list only one record for southern Africa, of an adult, presumably the same individual, that returned to West Coast National Park, Western Cape, South Africa, for four consecutive summers in 2000–2003. There are two subsequent records from the southern Africa sub-region, both from the Barra Peninsula, near Inhambane, Mozambique: a single in December 2004 and three individuals in September 2008 (M. Booysen *in Bull. ABC* 16: 107).



**Figures 1–2.** Great Knot *Calidris tenuirostris* with Bar-tailed Godwits *Limosa lapponica*, Curlew Sandpipers *C. ferruginea* and Sanderlings *C. alba*, Walvis Bay, Namibia, 24 November 2008 (Josh Engel)

Bécasseau de l'Anadyr *Calidris tenuirostris* avec Barges rousses *Limosa lapponica*, Bécasseaux cocorlis *C. ferruginea* et Bécasseaux sanderlings *C. alba*, Walvis Bay, Namibie, 24 novembre 2008 (Josh Engel)

---

## Acknowledgements

Thanks to Trevor Hardaker for supplying southern African records of Great Knot and to Callan Cohen for comments.

## References

- Aspinall, S. J. 1994. A wintering or passage population of Great Knot *Calidris tenuirostris* in the Arabian Gulf? *Wader Study Group Bull.* 72: 43–47.
- Cohen, C. & Winter, D. 2003. Great Knot *Calidris tenuirostris*: a new record for sub-Saharan Africa. *Bull. ABC* 10: 120–121.
- Green, M., McGrady, M., Newton, S. & Uttley, J. D. 1994. Counts of shorebirds at Barr al Hikman and Ghubbat al Hashish, Oman, winter 1989–90. *Wader Study Group Bull.* 72: 39–43.
- Hayman, P., Marchant, J. & Prater, T. 1986. *Shorebirds: An Identification Guide to the Waders of the World*. Boston: Houghton Mifflin.
- Hockey, P. A. R., Dean, W. R. J. & Ryan, P. G. (eds.) 2005. *Roberts' Birds of Southern Africa*. Seventh edn. Cape Town: Trustees of the John Voelcker Bird Book Fund.
- Lister, S. M. 1981. Le Grand Maubèche *Calidris tenuirostris* nouveau pour l'ouest du Paléarctique. *Alauda* 49: 227–228.
- Tropical Birding*, 1 Toucan Way, Bloubergrise 7441, South Africa; and 2653 Broadway, Evanston, IL 60201, USA. E-mail: jenge15230@yahoo.com

Received 24 March 2009; revision accepted 29 April 2009.